

Somatic Therapy Exercises For Trauma

Understanding Somatic Therapy and Its Role in Trauma Healing

Trauma is not just a psychological wound; it is a physiological event that leaves a lasting imprint on the body. When a person experiences a traumatic event, the nervous system can become stuck in a state of hyperarousal (fight or flight) or hypoarousal (freeze or collapse). Traditional talk therapy often addresses the cognitive and emotional aspects of trauma, but it may not fully reach the somatic, or body-based, layers of experience. This is where **somatic therapy exercises for trauma** come into play. These exercises are designed to help individuals reconnect with their bodily sensations, release stored tension, and restore a sense of safety and regulation.

Somatic therapy is a holistic approach that recognizes the mind-body connection as central to healing. By gently guiding attention to internal physical experiences, these exercises help trauma survivors process unresolved stress without having to verbally relive painful events. In this article, we will explore what somatic therapy is, why it is effective for trauma, and provide a comprehensive guide to specific **somatic therapy exercises for trauma** that you can incorporate into your healing journey. Whether you are new to this practice or looking to deepen your understanding, these techniques offer a path toward greater resilience and self-awareness.

What Is Somatic Therapy?

Somatic therapy is a body-centered therapeutic approach that focuses on the connection between the mind and the body. Developed by pioneers such as Peter Levine (Somatic Experiencing) and Pat Ogden (Sensorimotor Psychotherapy), it is grounded in the idea that trauma disrupts the autonomic nervous system's ability to self-regulate. Unlike purely cognitive therapies, somatic therapy uses physical awareness and movement to release trapped survival energy and restore equilibrium.

The term "somatic" comes from the Greek word *soma*, meaning "living body." In practice, somatic therapists help clients track bodily sensations—like tightness, heat, tingling, or numbness—and use these cues to process unresolved stress. For trauma survivors, this approach can be particularly effective because it bypasses the verbal parts of the brain that may shut down during traumatic recall. Instead, healing happens through gradual, titrated exposure to physical sensations in a safe and controlled manner.

How Trauma Affects the Body

When faced with a threat, the body responds instinctively. The sympathetic nervous system activates, flooding the body with stress hormones like cortisol and adrenaline to prepare for fight or flight. If neither option is possible, the

parasympathetic system initiates a freeze response. In a healthy situation, after the danger passes, the nervous system returns to a baseline state of calm. But for trauma survivors, this cycle can become incomplete. The energy that would have been used for action remains locked in the body, leading to chronic tension, hypervigilance, anxiety, or numbness.

These physical manifestations often go unnoticed or are dismissed. People may experience chronic pain, digestive issues, headaches, or a sense of being disconnected from their own body. **Somatic therapy exercises for trauma** aim to complete these thwarted responses by allowing the body to gently discharge stored energy. Through mindful awareness and small, manageable movements, the nervous system learns that it is safe again.

Key Principles of Somatic Exercises for Trauma

Before diving into specific exercises, it is helpful to understand the core principles that make somatic work safe and effective for trauma survivors:

- **Pendulation:** The practice of gently moving between sensations of discomfort or activation and areas of ease, safety, or neutrality in the body. This rhythm helps the nervous system expand its capacity to handle distress without becoming overwhelmed.
- **Titration:** Gradual exposure to small amounts of traumatic material, avoiding full reactivation. Titration respects the individual's window of tolerance—the zone where they can feel sensations without going into extreme hyper- or hypoarousal.
- **Resourcing:** Building inner strengths by identifying sensations, memories, or images that evoke a sense of calm, safety, or support. A

resource could be a pleasant memory, a beloved pet, or a feeling of warmth in the hands.

- **Grounding:** Bringing attention into the present moment through physical contact with the earth or supportive surfaces. Grounding helps counteract dissociation and creates a foundation for exploring sensations.
- **Self-compassion:** Approaching the process with kindness and patience. Trauma work can bring up vulnerability, and it is essential to move at your own pace.

Effective Somatic Therapy Exercises for Trauma

Below are a series of **somatic therapy exercises for trauma** that you can practice on your own or with a qualified therapist. Always listen to your body's signals. If an exercise feels overwhelming, stop and return to a grounding resource. These exercises are designed to be gentle and incremental.

1. Grounding Through the Soles of the Feet

This foundational exercise helps bring your awareness out of your head and into your body, particularly the contact point with the floor. It is excellent for moments of anxiety, flashbacks, or dissociation.

1. Stand or sit comfortably with your feet flat on the ground. If standing, keep your knees slightly soft.
2. Close your eyes or soften your gaze. Take a few natural breaths.
3. Bring your attention to the soles of your feet. Notice the pressure against the floor—is it even? Is one foot heavier than the other?

Somatic Therapy Exercises For Trauma

4. Slowly shift your weight slightly forward and back, then side to side. Feel how the fabric of your shoes or socks touches your skin. If barefoot, feel the texture of the floor.
5. Now imagine roots gently growing from the soles of your feet into the earth. Breathe calmly and stay with this sensation for 1–2 minutes.
6. When ready, slowly take a deeper breath and notice any changes in your body.

Why it works: Grounding interrupts the dissociative response and anchors you in the present. The physical focus on the feet activates proprioception, which can calm the amygdala and reduce hypervigilance.

2. Resourcing: Finding Your Safe Place Within

Resourcing is a crucial step before exploring any trauma-related sensations. It builds an internal reservoir of safety that you can draw upon when needed.

1. Sit in a comfortable position and bring your hands to rest on your thighs or heart.
2. Think of a person, place, memory, or even an object that makes you feel safe or loved. It could be a cozy room, a childhood pet, a kind friend, or a place in nature.
3. Allow an image or feeling to arise. Notice where in your body you sense this resource—perhaps a warmth in your chest, a gentle relaxation in your shoulders, or lightness in your belly.
4. Stay with this sensation for 15–30 seconds. Let it grow. You can gently touch the area where you feel it (e.g., placing a hand on your heart).
5. If the feeling fades, return to the image and breathe. Over time, you can access this resource with a single breath or touch.

Why it works: Positive resources regulate the nervous system by activating

the parasympathetic branch. They create a "anchor" of safety that can be used during more challenging exercises.

3. Pendulation Between Discomfort and Ease

Pendulation is a core skill in somatic therapy. It prevents retraumatization by maintaining dual awareness—feeling both the activation and a safe sensation.

1. Begin by identifying a minor tightness, ache, or discomfort in your body. It could be a knot in your shoulder, a slight heaviness in your chest, or tension in your jaw.
2. Place your attention there. Notice the quality: is it sharp, dull, hot, cold, pulsing? Stay for just a few seconds.
3. Now deliberately shift your attention to a part of your body that feels neutral or pleasant—perhaps your warm hands, the gentle rise and fall of your belly, or the resource you cultivated earlier.
4. Stay with the ease for longer than you stayed with the discomfort (e.g., 10 seconds of discomfort, then 20 seconds of ease).
5. Gently oscillate back and forth a few times. You may notice the discomfort soften or change.

Why it works: Pendulation teaches the nervous system that discomfort is temporary and that safety is always available. This builds tolerance and reduces fear of internal sensations.

4. Body Scanning with Gentle Curiosity

A body scan can help you uncover hidden areas of tension or numbness without judgment. For trauma survivors, it is important to keep the scan brief and to anchor with touch if needed.

1. Lie down or sit comfortably. Close your eyes and take a few breaths.
2. Begin at the top of your head. Simply notice—what do you feel? Itching, pressure, nothing at all? Don't try to change anything.
3. Slowly move your attention down your face, neck, shoulders, arms, hands, torso, hips, legs, and feet.
4. If you encounter an area of intense sensation, stop and use pendulation: shift your attention to a neutral spot (like your left little finger) for a few seconds, then gently return to the intense area. You can also place a hand on that spot for comfort.
5. If you feel completely numb in a region, acknowledge that numbness by saying to yourself, "This part feels numb. That's okay."
6. Complete the scan by returning to an area of safety—perhaps the breath or the hands.

Why it works: A trauma-informed body scan reduces avoidance of physical experience. It helps you map where your nervous system holds activation or shut-down, which is the first step toward releasing it.

5. Completing the Fight Response in Slow Motion

This exercise allows the body to discharge the incomplete "fight" energy in a safe, controlled way. It should only be done when you have robust resources and are in a stable state.

1. Stand with feet hip-width apart, knees bent slightly. Imagine a situation that triggers mild frustration (not a major trauma).
2. Notice any impulse in your arms or hands to push, squeeze, or hit. You may feel a subtle tension in your fists or shoulders.
3. Now, very slowly—as if underwater—start to extend your arms in front of you, as though you are pushing a heavy object. Go as slowly as

possible.

4. Make a sound if it feels natural, like a soft "ha" or a grunt. Keep your jaw relaxed.
5. Continue pushing slowly until your arms are fully extended. Then pause, take a breath, and even more slowly, bring your arms back to your sides.
6. Notice if the initial tension has lessened. You may feel a vibration, warmth, or relaxation.

Why it works: This micro-movement allows the body to complete a thwarted motor impulse safely, releasing the stored energy without overwhelming the system. The slowness keeps the experience within the window of tolerance.

6. Orienting: A Return to the Present Environment

Trauma often leaves us scanning for threats in a "reptilian" way. Orienting exercises help the nervous system recognize that the current environment is safe.

1. Pause and turn your head slowly to the right. Notice what you see, hear, and smell. Take in the full corner of the room.
2. Turn your head slowly to the left. Again, notice details—colors, objects, sounds.
3. Look up toward the ceiling, then down toward the floor. Let your eyes move naturally.
4. Now choose one object in the room. Look at it with soft focus, noticing its shape, texture, and color. Allow your peripheral vision to expand.
5. Take a breath and feel your feet on the ground.

Why it works: Orienting engages the social engagement system (ventral vagal pathway), which is responsible for calm connection. It counters the hyper